

# verno<sup>®</sup>

## CaB

### Micronutrient Fertilizer

**Boron (B)** 90 g/kg (9%)  
**Calcium (Ca)** 160 g/kg (16%)

**Water dispersible & dust-free granules for the correction of micronutrient deficiencies.**

**VERNO CaB** is a micronutrient fertilizer composed of micronized particles that strongly adhere to plant surfaces. The particles act as reservoirs for nutrients with slow release providing mineral nutrition over time. To be used only where there is a recognized need. Do not exceed the recommended application rate.

#### INSTRUCTIONS FOR USE:

Cereals	1 - 2 kg/ha from 2-leaf until first node visible (growth stage 31)
Oilseed Rape	1 - 3 kg/ha at 4-9 leaf stage (BBCH 14-19, stem elongation (BBCH 30), and petal fall (BBCH 67)
Potatoes	1 - 2 kg/ha about 14 days after full emergence (BBCH 09), repeat after 14 days
Fruit Trees	1 - 3 kg/ha at pink bud, the beginning of flowering, and petal fall
Sugar Beets	1 - 2 kg/ha at 4-6 leaf stage, repeat after 14 days if necessary
Soy/Legumes	1 - 2 kg/ha at 4-6 leaf stage, repeat every 14 days if necessary up to one month prior to harvest
Citrus	1 - 3 kg/ha before flowering and immediately after flowering
Coffee	1 - 3 kg/ha before flowering and immediately after flowering
Cherry	1 - 3 kg/ha from flower bud through flowering, and early post harvest

#### APPLICATION GUIDELINES:

Fill the tank  $\frac{3}{4}$  full with water and add the correct dose with agitation. Maintain agitation during application. Use 200-1000 liters of water per hectare, depending on spray equipment. Multiple low-rate applications are recommended.

#### MIXABILITY WITH OTHER PRODUCTS:

VERNO CaB is mixable with all other VERNO formulations. Tank mixes are usually not recommended unless mixability is confirmed by the manufacturer of the mixing partner. Mixing is not recommended when the pH of the spray liquid is lower than pH 6. If you decide to make a mixture with other products, VERNO CaB should be suspended in water before adding the mixing partner. Look for precipitate that can clog spray equipment and avoid using the mixture if this occurs. It is recommended that you perform a test by spraying a limited area of crop to assess plant tolerance as a general precaution. Crop damage/phytotoxicity is usually visible within 2 days.

#### STORAGE

Shelf life is 3 years from the date of manufacture if the product is stored in the closed original packaging and kept dry. Tolerates freezing.

#### PRECAUTIONS:

Keep out of reach of children. Wear gloves, goggles, and protective clothing when handling and spraying. Dispose of empty and clean packaging in accordance with local regulations. Avoid contamination of water bodies during spraying. If tank is not empty following treatment, use a 10-fold dilution and spray over the treated crop.

Manufactured by:  
NORDOX AS  
Øststensjøveien 13, 0661  
Oslo, Norway  
Tel: +47 22975000  
www.nordox.no



Made in Norway



Net contents:  
**8 kg**